

Generalized Bicircular Projections on Some Matrix and Operator Spaces

Dijana Ilišević

University of Zagreb, Croatia

Applied Linear Algebra (ALA 2010)
Novi Sad, Serbia, May 24–28, 2010

Definition of a Bicircular Projection

Definition

Let \mathcal{X} be a complex Banach space and let $P : \mathcal{X} \rightarrow \mathcal{X}$ be a linear projection. A projection P is called **bicircular** if the mapping $P + \lambda \bar{P}$ is an isometry for all modulus one complex numbers λ .

Example

Every orthogonal projection on a complex Hilbert space is bicircular.

Symmetric and Antisymmetric Operators

Let $B(\mathcal{H})$ be the algebra of all bounded linear operators on a complex Hilbert space \mathcal{H} . Throughout we fix an orthonormal basis $\{e_\lambda : \lambda \in \Lambda\}$ of \mathcal{H} .

Let $T \in B(\mathcal{H})$. If $S \in B(\mathcal{H})$ is such that

$$\langle Te_\lambda, e_\mu \rangle = \langle Se_\mu, e_\lambda \rangle \quad (\lambda, \mu \in \Lambda),$$

then S is called **the transpose of T associated to the basis $\{e_\lambda : \lambda \in \Lambda\}$** and it is denoted by T^t .

- $S(\mathcal{H}) = \{T \in B(\mathcal{H}) : T^t = T\}$ **symmetric operators**
- $A(\mathcal{H}) = \{T \in B(\mathcal{H}) : T^t = -T\}$ **antisymmetric operators**

Bicircular Projections on Some Operator Spaces

Theorem (L.L. Stachó and B. Zalar, LAA, 2004)

- (i) Let $P : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ be a bicircular projection. Then P has the form $X \mapsto QX$ or $X \mapsto XQ$ for some $Q = Q^* = Q^2 \in B(\mathcal{H})$.
- (ii) Let $P : S(\mathcal{H}) \rightarrow S(\mathcal{H})$ be a bicircular projection. Then either $P = 0$ or $P = I$.
- (iii) Let $P : A(\mathcal{H}) \rightarrow A(\mathcal{H})$ be a bicircular projection. Then P or \overline{P} has the form $X \mapsto QX + XQ^t$ with $Q = x \otimes x$ for some unit vector $x \in \mathcal{H}$.

- the structure of bicircular projections on an arbitrary C^* -algebra:
M. Fošner and D. Ilišević, Comm. Algebra, 2005.

Bicircular Projections on Some Operator Spaces

Theorem (L.L. Stachó and B. Zalar, LAA, 2004)

- (i) Let $P : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ be a bicircular projection. Then P has the form $X \mapsto QX$ or $X \mapsto XQ$ for some $Q = Q^* = Q^2 \in B(\mathcal{H})$.
- (ii) Let $P : S(\mathcal{H}) \rightarrow S(\mathcal{H})$ be a bicircular projection. Then either $P = 0$ or $P = I$.
- (iii) Let $P : A(\mathcal{H}) \rightarrow A(\mathcal{H})$ be a bicircular projection. Then P or \overline{P} has the form $X \mapsto QX + XQ^t$ with $Q = x \otimes x$ for some unit vector $x \in \mathcal{H}$.

- the structure of bicircular projections on an arbitrary C^* -algebra:
M. Fošner and D. Ilišević, Comm. Algebra, 2005.

Hermitian Projections

Definition

A bounded linear operator $T : \mathcal{X} \rightarrow \mathcal{X}$ is said to be **hermitian** if $e^{i\varphi T}$ is an isometry for all $\varphi \in \mathbb{R}$.

Theorem (J. Jamison, LAA, 2007)

A linear projection on \mathcal{X} is a bicircular projection if and only if it is a hermitian projection.

Definition of a Generalized Bicircular Projection

Definition

A projection $P : \mathcal{X} \rightarrow \mathcal{X}$ is called **generalized bicircular** if the mapping $P + \lambda \overline{P}$ is an isometry for some modulus one complex number $\lambda \neq 1$.

- These mappings were first studied by M. Fošner, D. Ilišević and C.K. Li, LAA, 2007.
- The term “generalized bicircular projection” (GBP) first appeared in a paper by F. Botelho and J. Jamison, PAMS, 2008.

GBP on $S_n(\mathbb{C})$

Let A be $S_n(\mathbb{C})$ or $K_n(\mathbb{C})$. A norm $\|\cdot\|$ on A is said to be a **unitary congruence invariant norm** if

$$\|UXU^t\| = \|X\|$$

for all unitary $U \in M_n(\mathbb{C})$ and all $X \in A$.

Theorem (M. Fošner, D. Ilišević and C.K. Li, LAA, 2007)

Let $\|\cdot\|$ be a unitary congruence invariant norm on $S_n(\mathbb{C})$, which is not a multiple of the Frobenius norm, and let \mathcal{K} be the isometry group of $\|\cdot\|$. Suppose $P : S_n(\mathbb{C}) \rightarrow S_n(\mathbb{C})$ is a non-trivial linear projection and $\lambda \neq 1$ a modulus one complex number. Then $P + \lambda\bar{P} \in \mathcal{K}$ if and only if $\lambda = -1$ and there exists $Q = Q^ = Q^2 \in M_n(\mathbb{C})$ such that P or \bar{P} has the form $X \mapsto QXQ^t + (I - Q)X(I - Q^t)$.*

GBP on $S_n(\mathbb{C})$

Let A be $S_n(\mathbb{C})$ or $K_n(\mathbb{C})$. A norm $\|\cdot\|$ on A is said to be a **unitary congruence invariant norm** if

$$\|UXU^t\| = \|X\|$$

for all unitary $U \in M_n(\mathbb{C})$ and all $X \in A$.

Theorem (M. Fošner, D. Ilišević and C.K. Li, LAA, 2007)

Let $\|\cdot\|$ be a unitary congruence invariant norm on $S_n(\mathbb{C})$, which is not a multiple of the Frobenius norm, and let \mathcal{K} be the isometry group of $\|\cdot\|$. Suppose $P : S_n(\mathbb{C}) \rightarrow S_n(\mathbb{C})$ is a non-trivial linear projection and $\lambda \neq 1$ a modulus one complex number. Then $P + \lambda\bar{P} \in \mathcal{K}$ if and only if $\lambda = -1$ and there exists $Q = Q^ = Q^2 \in M_n(\mathbb{C})$ such that P or \bar{P} has the form $X \mapsto QXQ^t + (I - Q)X(I - Q^t)$.*

GBP on $K_n(\mathbb{C})$ **Theorem (M. Fošner, D. Ilišević and C.K. Li, LAA, 2007)**

Let $n \geq 3$ and $\|\cdot\|$ be a unitary congruence invariant norm on $K_n(\mathbb{C})$, which is not a multiple of the Frobenius norm. Let \mathcal{K} be the isometry group of $\|\cdot\|$. Suppose $P : K_n(\mathbb{C}) \rightarrow K_n(\mathbb{C})$ is a non-trivial linear projection and $\lambda \neq 1$ a modulus one complex number. Then $P + \lambda\bar{P} \in \mathcal{K}$ if and only if one of the following holds.

- (a) There exists $Q = vv^*$ for a unit vector $v \in \mathbb{C}^n$ such that P or \bar{P} has the form $X \mapsto QX + XQ^t$.
- (b) $\lambda = -1$, $\mathcal{K} = G$ and there exists $Q = Q^* = Q^2 \in M_n(\mathbb{C})$ such that P or \bar{P} has the form $X \mapsto QXQ^t + (I - Q)X(I - Q^t)$.
- (c) $(\lambda, n) = (-1, 4)$, $\psi \in \mathcal{K}$, and there is $U \in U(\mathbb{C}^4)$, satisfying $\psi(U^t X U) = \bar{U}\psi(X)U^*$ for all $X \in K_4(\mathbb{C})$, such that P or \bar{P} has the form $X \mapsto (X + \psi(U^t X U))/2 = (X + \bar{U}\psi(X)U^*)/2$.

JB*-triples

A **JB*-triple** is a complex Banach space A together with a continuous triple product $\{\cdot \cdot \cdot\} : A \times A \times A \rightarrow A$ such that

- (i) $\{xyz\}$ is linear in x and z and conjugate linear in y ;
- (ii) $\{xyz\} = \{zyx\}$;
- (iii) for any $x \in A$, the operator $\delta(x) : A \rightarrow A$ defined by $\delta(x)y = \{xxy\}$ is hermitian with nonnegative spectrum;
- (iv) $\delta(x)\{abc\} = \{\delta(x)a, b, c\} - \{a, \delta(x)b, c\} + \{a, b, \delta(x)c\}$;
- (v) for every $x \in A$, $\|\{xxx\}\| = \|x\|^3$.

- complex Hilbert spaces: $\{xyz\} = \frac{1}{2}(\langle x, y \rangle z + \langle z, y \rangle x)$
- C*-algebras, $S(\mathcal{H})$, $A(\mathcal{H})$: $\{xyz\} = \frac{1}{2}(xy^*z + zy^*x)$.

GBP on JB^* -triples**Theorem (D. Ilišević, LAA, 2010)**

Let A be a JB^* -triple and let $P : A \rightarrow A$ be a linear projection. Then $P + \lambda \bar{P}$ is an isometry for some modulus one complex number $\lambda \neq 1$ if and only if one of the following holds:

- (i) P is hermitian (\equiv bicircular),
- (ii) $\lambda = -1$ and $P = \frac{1}{2}(I + \varphi)$ for some linear isometry $\varphi : A \rightarrow A$ satisfying $\varphi^2 = I$.

GBP on Arbitrary Complex Banach Spaces

Theorem (P.-K. Lin, JMAA, 2008)

Let \mathcal{X} be a complex Banach space and let $P : \mathcal{X} \rightarrow \mathcal{X}$ be a linear projection. Then $P + \lambda\bar{P}$ is an isometry for some modulus one complex number $\lambda \neq 1$ if and only if one of the following holds:

- (i) P is hermitian (\equiv bicircular),
- (ii) $\lambda = e^{\frac{2\pi i}{n}}$ for some integer $n \geq 2$.

Furthermore, if n is any integer such that $n \geq 2$, then for $\lambda = e^{\frac{2\pi i}{n}}$ there is a complex Banach space \mathcal{X} and a nontrivial linear projection P on \mathcal{X} such that $P + \lambda\bar{P}$ is an isometry.

GBP on Some Matrix and Operator Spaces

- bicircular projections on $S(\mathcal{H})$ and $A(\mathcal{H})$:
L.L. Stachó and B. Zalar, LAA, 2004
- generalized bicircular projections on $S_n(\mathbb{C})$ and $K_n(\mathbb{C})$:
M. Fošner, D. Ilišević and C.K. Li, LAA, 2007
- generalized bicircular projections on $S(\mathcal{H})$ and $A(\mathcal{H})$:
A. Fošner and D. Ilišević, submitted

GBP on Some Matrix and Operator Spaces

- bicircular projections on $S(\mathcal{H})$ and $A(\mathcal{H})$:
L.L. Stachó and B. Zalar, LAA, 2004
- generalized bicircular projections on $S_n(\mathbb{C})$ and $K_n(\mathbb{C})$:
M. Fošner, D. Ilišević and C.K. Li, LAA, 2007
- generalized bicircular projections on $S(\mathcal{H})$ and $A(\mathcal{H})$:
A. Fošner and D. Ilišević, submitted

Surjective Linear Isometries on $S(\mathcal{H})$ and $A(\mathcal{H})$

Every surjective linear isometry $\varphi : A \rightarrow A$, where A is $S(\mathcal{H})$ or $A(\mathcal{H})$, satisfies

$$\varphi(XY^*X) = \varphi(X)\varphi(Y)^*\varphi(X)$$

for all $X, Y \in A$.

The following theorem gives an explicit formula for φ .

Theorem (A. Fošner and D. Ilišević, submitted)

Let A be $S(\mathcal{H})$ or $A(\mathcal{H})$ and let $\varphi : A \rightarrow A$ be a surjective linear isometry. Then there exists a unitary $U \in B(\mathcal{H})$ such that φ has the form $X \mapsto UXU^t$.

GBP on $S(\mathcal{H})$ and $A(\mathcal{H})$

Corollary

Let $P : S(\mathcal{H}) \rightarrow S(\mathcal{H})$ be a nontrivial linear projection and $\lambda \neq 1$ a modulus one complex number. Then $P + \lambda\bar{P}$ is an isometry if and only if $\lambda = -1$ and there exists $Q = Q^* = Q^2 \in B(\mathcal{H})$ such that P or \bar{P} has the form $X \mapsto QXQ^t + (I - Q)X(I - Q^t)$.

Corollary

Let $P : A(\mathcal{H}) \rightarrow A(\mathcal{H})$ be a nontrivial linear projection and $\lambda \neq 1$ a modulus one complex number. Then $P + \lambda\bar{P}$ is an isometry if and only if one of the following holds:

- (i) P or \bar{P} has the form $X \mapsto QX + XQ^t$, where $Q = x \otimes x$ for some norm one $x \in \mathcal{H}$,
- (ii) $\lambda = -1$ and there exists $Q = Q^* = Q^2 \in B(\mathcal{H})$ such that P or \bar{P} has the form $X \mapsto QXQ^t + (I - Q)X(I - Q^t)$.